

தமிழ்நாடு தேர்வுத்துறை / இலங்கைப் பரீட்சைத் திணைக்களம் / Department of Examinations,
சென்னை / பொதுத் தராதரப்பத்திர(உயர் தர)ப் பரீட்சை, 1999 ஓகஸ்ட்
General Certificate of Education (Adv. Level) Examination, August 1999

உயர் தர பரீட்சை II

இரசாயனவியல் II

Chemistry II

மூன்று மணி / மூன்று மணித்தியாலங்கள் / Three hours

02

E | II

Index No.:

Important : This question paper consists of **four** sheets. Put the sheets together in the correct **order of pages** before answering.

Use of calculators is not allowed.

This question paper consists of three parts A, B and C. The time allotted for all three parts is three hours.

PART A — STRUCTURED ESSAY

Answer all the questions. Write your answer in the space provided below each question. Please ensure that the space provided is sufficient for the answer and that extensive answers are not expected.

PART B AND PART C—ESSAY

Answer four questions selecting two questions from each part. Use the paper supplied for this part. At the end of the time allotted for this paper, tie the three parts A, B and C together so that Part A is on top and hand them over to the supervisor. You are permitted to remove only Parts B and C of the question paper from the Examination Hall.

Universal gas constant (R) = $8.314 \text{ J K}^{-1} \text{ mol}^{-1}$

N.B. The following abbreviations have been used.

aq = aqueous

C = Celsius or Centigrade or Coulomb

g = gas or gram

l = liquid

mol dm^{-3} = moles per cubic decimetre

s = solid or second

Other abbreviations also follow standard usage.

PART A — STRUCTURED ESSAY

Answer all four questions. Each question carries 10 marks.

1. (a) (i) Define the term, 'atomic mass unit' (amu).

Do not
write
anything
in this
column

(ii) It is possible that the mass of a certain molecule of carbon dioxide supplied to you is **not** equal to 44.0 amu. Give **two** reasons for this possibility.

(b) (i) The element, X belongs to the 4th Period of the Periodic Table. An atom of X forms an anion. The highest valency-state oxide formed by X is XO_3 . Identify X.

(ii) The compound, XCl_4 formed by the above element X is reduced with LiAlH_4 . Write the molecular formula of the hydride of X that is most likely to be formed in this reaction.

not
e
hing
is
mn

(c) Can the hydride referred to in (b)(ii) above, act as an oxidizing agent? If so, give reasons for that.

2. (a) Assume that $\text{Na}_2\text{S}_2\text{O}_3$ reacts with HNO_3 under a certain condition forming NO and water only. Write a balanced chemical equation for this reaction, either by considering oxidation numbers or by another method.

Do not
write
anything
in this
column

- (b) A finely powdered mixture has been prepared so that the $\text{CaCO}_3 : \text{MgCO}_3$ mole ratio is $1 : x$. It contains no other substances. 1.30 g of this mixture was heated strongly, until it was converted completely into CaO and MgO. The mass of the mixture of oxides thus obtained was 0.640 g.

The relative atomic masses relevant here are as follows:

$$\text{Ca} = 40.0; \text{Mg} = 24.0; \text{O} = 16.0; \text{C} = 12.0.$$

- (i) Now consider P , Q , R and S in the mathematical expression given below.

$$\frac{100 + P \times x}{Q + R \times x} = \frac{1.30 \text{ g}}{S}$$

Write the values appropriate for P , Q , R and S in the Table provided below.


P	
Q	
R	
S	

- (ii) Substitute these values appropriately in the mathematical expression given above, and calculate the value of x .

- (c) (i) State Gay Lussac's Law.

- (ii) Direct your attention now to the experiment that you carried out in the laboratory for the purpose of demonstrating Gay Lussac's Law, using equal volumes of hydrogen gas and chlorine gas under constant temperature and pressure.

Assume that you are supplied with a sample of hydrogen gas and a sample of chlorine gas as shown below.





Describe briefly how you would demonstrate in the laboratory that Gay Lussac's Law is true, using the two gas samples shown above.

3. (a) The organic compound, Y contains C, H and O only. When Y is subjected to complete combustion, CO_2 and H_2O are obtained in the mole ratio of 2 : 1, respectively. The accurate relative molecular mass of Y is equal to 152. The percentage of O in Y is less than 40%, by weight. Determine the molecular formula of Y.



The relevant relative atomic masses are as follows: C = 12.0; H = 1.00; O = 16.0

- (b) A and B are two alkynes. A and B were reacted as shown below, and C and D were obtained, respectively.



The molecular formulae of both C and D were C_4H_8 .

C and D reacted as follows:



Do not
write
anything
in this
column


(i) You are supplied with a mixture containing both A and B. How would you attempt to obtain either pure A or pure B from this mixture by a chemical method?

(ii) Draw the structures possible for the compounds C and D.





Do not write anything in this column

- (iii) Draw the structure of one of the optical isomers formed in the reaction with HBr, in the usual manner, so that its three-dimensional nature is clearly apparent.

- (c) Consider the following series of reactions.



- (i) Draw appropriately the structures of E, F, G and H in the cages provided below.



- (ii) Identify L and M and appropriately indicate them in the cages provided below.

L	
M	


ot

ing

s


nn

(d) Assume that you are supplied with



as the **only** organic compound.


Indicate how you would synthesize



using the above compound.

N.B. Disregard the isomerism of the products. If the method of synthesis proposed is unnecessarily long, you will not be awarded the maximum marks.

1. (a) Name the compound with the following structure, in accordance with IUPAC nomenclature.



Do not
write
anything
in this
column

- (c) (i) A brief description pertaining to the reaction between aqueous KCN and the bromoalkane, $R_1R_2R_3CBr$ is given below.

"This reaction is called $\left\{ \begin{array}{l} \text{a free radical /} \\ \text{an electrophillic /} \\ \text{a nucleophillic} \end{array} \right\} \left\{ \begin{array}{l} \text{substitution /} \\ \text{addition} \end{array} \right\}$ reaction."


Clearly cross out the inappropriate words out of the five words within the brackets.

N.B. The appropriate words should be clearly visible. If your responses are not clear, you will not be awarded marks.

- (ii) Present clearly the mechanism of the reaction given in 4 (c)(i) above.

- (d) Propose a method for effecting the following conversion.

N.B. If the method of conversion proposed by you is unnecessarily long, you will not be awarded the maximum marks.




it

ing


n

(d) Assume that you are supplied with



as the only organic compound.


Indicate how you would synthesize



using the above compound.

N.B. Disregard the isomerism of the products. If the method of synthesis proposed is unnecessarily long, you will not be awarded the maximum marks.

4. (a) Name the compound with the following structure, in accordance with IUPAC nomenclature.



ශ්‍රී ලංකා විභාග දෙපාර්තමේන්තුව / இலங்கைப் பரீட்சைத் திணைக்களம் / Department of Examinations, Sri Lanka

අධ්‍යයන පොදු සහතික පත්‍ර (උසස් පෙළ) විභාගය, 1999 අගෝස්තු
கல்விப் பொதுத் தராதரப்பத்திர(உயர் தர)ப் பரீட்சை, 1999 ஓகஸ்த்
General Certificate of Education (Adv. Level) Examination, August 1999

රසායන විද්‍යාව II
இரசாயனவியல் II
Chemistry II

02
E | II

PART B — ESSAY

Answer two questions only. Each question carries 15 marks.

5. (a) Using the equation, $PV = \frac{1}{3}mNc^2$ pertaining to the Kinetic Theory of gases, derive Dalton's Law of partial pressures.
- (b) A mixture of gases contains 75.0% of N_2 gas and 25.0% of O_2 gas, by volume. The pressure of the mixture of gases is $1.00 \times 10^5 \text{ N m}^{-2}$ and the temperature is 300 K. Assuming ideal behaviour calculate the following:
- The partial pressure of O_2 in this mixture of gases.
 - The relative molecular mass relevant for this mixture of gases. (The relative atomic masses of N and O are 14.0 and 16.0 respectively.)
 - The density of this mixture of gases.
- (c) You are supplied with a real gas. You are not informed of its relative molecular mass. Explain how you would attempt to show that this real gas does not behave ideally.
- (d) You would have performed an experiment in the laboratory for the purpose of determining the molar volume of oxygen gas. Describe that experiment briefly, and explain how the molar volume of O_2 at s.t.p. is determined.
6. (a) At a certain temperature, the K_c value for the equilibrium, $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ is about 25. Assume that this reaction was started with equi-molar amounts of $H_2(g)$ and $I_2(g)$. Draw a diagram of the graphs which shows the variation of $H_2(g)$ concentration and $HI(g)$ concentration with time, for this reaction at the above temperature.
- (b) 0.200 mol NO, 0.100 mol H_2 and 0.200 mol H_2O were placed initially in a closed vessel of volume 0.0200 m^3 . At 500 K the following equilibrium occurred.
- $$2NO(g) + 2H_2(g) \rightleftharpoons N_2(g) + 2H_2O(g)$$
- At equilibrium, the amount of NO(g) present was 0.150 mol.
- Calculate the K_c for this equilibrium.
 - Using the value obtained above for K_c , calculate the K_p for this equilibrium.
- (c) (i) Derive the relationship between K_b of NH_3 and K_a of NH_4^+ , in aqueous solution.
- (ii) At 25 °C, the concentration of an aqueous solution of a mono-basic weak acid is 0.10 mol dm^{-3} . Calculate the OH^- concentration in this aqueous solution.

$$K_a \text{ at } 25 \text{ }^\circ\text{C} = 9.0 \times 10^{-9} \text{ mol dm}^{-3}$$

$$K_w \text{ at } 25 \text{ }^\circ\text{C} = 1.0 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$$

95

[Turn over

- (d) The ionic compound, FeX_2 is only slightly soluble in water. The anion, X^- is not oxidized in a solution by chemical means. The amount of FeX_2 dissolving in water cannot be directly determined by methods involving weighings. However, there is a measurable Fe^{2+} ion concentration in a saturated aqueous solution at 25 °C. Explain how you would attempt to determine the solubility product of FeX_2 at 25 °C, under these conditions.

7. (a) (i) Consider the following data:

The standard enthalpy of formation of $\text{CH}_4(\text{g}) = -75.0 \text{ kJ mol}^{-1}$

The standard enthalpy change for the reaction, $2\text{H}(\text{g}) \rightarrow \text{H}_2(\text{g}) = -432 \text{ kJ mol}^{-1}$

The standard enthalpy change for the reaction, $\text{C}(\text{graphite}) \rightarrow \text{C}(\text{g}) = +715 \text{ kJ mol}^{-1}$


Using these data, calculate the standard mean bond dissociation enthalpy of C-H pertaining to $\text{CH}_4(\text{g})$.

- (ii) Three standard bond dissociation enthalpies are given below.

bond	standard bond dissociation enthalpy
H — H	+ 432 kJ mol ⁻¹
F — F	+ 158 kJ mol ⁻¹
H — F	+ 569 kJ mol ⁻¹

Using these values, calculate the standard enthalpy of formation of $\text{HF}(\text{g})$.

- (b) (i) Describe briefly what is meant by 'the standard hydrogen electrode'.
 (ii) Describe briefly what is meant by 'the standard zinc electrode'.
 (c) (i) A certain electrochemical cell which is at 25 °C is represented below.




Using the data given below, calculate the e.m.f. of the above cell.

E^\ominus values at 25 °C are as follows:

$$E_{\text{Sn}^{2+}/\text{Sn}}^\ominus = -0.136 \text{ V}; E_{\text{Cu}^{2+}/\text{Cu}}^\ominus = +0.337 \text{ V}$$

- (ii) At which electrode does the reduction take place when a current of electricity is drawn from the cell? Write the electrode reaction that takes place at that electrode.
 (d) (i) When the temperature increases slightly, the rate of a chemical reaction increases considerably. Explain this fact by considering the distribution of the velocities of the molecules (Boltzmann Curve).

- (ii) Assume that the solid organic compound,  dissolves in water. Assume also that one molecule of this compound, in aqueous solution, hydrolyses slowly giving two molecules of $\text{R}-\text{O}-\text{H}$ and another product, and also that the two molecules of $\text{R}-\text{O}-\text{H}$ are formed simultaneously. Describe briefly how you would attempt to determine the order of this reaction.

PART C — ESSAY

Answer two questions only. Each question carries 15 marks.

8. (a) (i) Write an expression which will describe in a generalized form the electronic configurations of all of the following elements: Be, Mg, Ca and Sr.
- (ii) Write an expression which will describe in a generalized form the electronic configurations of all of the following elements: F, Cl, Br and I.
- (b) (i) Predict what products would be formed when CuCl is reacted with caesium.
- (ii) Explain the above reaction on the basis of ionization energies.
- (c) (i) Write a balanced chemical equation for the reaction that occurs between H_2O_2 , KMnO_4 and an excess of dilute H_2SO_4 .
- (ii) How would you explain the action of H_2O_2 as a reducing agent under the above conditions?
- (d) An aqueous solution supplied to you contains the two compounds Na_2CO_3 and NaOH. You are also supplied with a standard $0.100 \text{ mol dm}^{-3}$ H_2SO_4 solution. However, you are not supplied with other acids or bases. As the indicator you are given only phenolphthalein. Explain how you would determine volumetrically the NaOH concentration present in the original solution supplied to you, under these conditions.

N.B. You are provided with the other materials and facilities available in the Chemistry Laboratory.

9. (a) This part of the question pertains to the industrial production of caustic soda.
- (i) Making use of a labelled diagram, indicate clearly the essential parts and all the relevant materials used in a diaphragm cell.
- (ii) What is the principal purpose served by the diaphragm?
- (iii) Write the chemical reaction that takes place at the anode.
- (iv) Write the chemical reaction that takes place at the cathode.
- (v) Write, in the balanced form, the overall chemical reaction that takes place in the formation of NaOH in the diaphragm cell.
- (b) Name the following compounds in accordance with IUPAC nomenclature.
- (i) $[\text{Cu}(\text{NH}_3)_4]\text{Br}_2$
- (ii) $(\text{NH}_4)_3[\text{Fe}(\text{CN})_6]$
- (c) Explain the following facts pertaining to the Periodic Table.
- (i) In comparison with the melting points of the elements in the *s*-block, the melting points of the elements in the *d*-block are very high.
- (ii) The elements manganese and bromine both belong to the same period in the Periodic Table. The total number of electrons in the two outermost sub-energy levels of the atoms of both these elements is equal to seven. However, manganese is a good conductor of electricity, while bromine does not conduct electricity.
- (d) You are supplied with several samples of a solution containing CuCl_2 , FeCl_3 and NiCl_2 . How would you demonstrate clearly that the cations, Cu^{2+} , Fe^{3+} and Ni^{2+} are present in this solution?

10. (a) Write a brief description of how the environment could be polluted by the industrial production of ammonia by the Haber Process.
N.B. It is sufficient to present five important facts.
- (b) (i) The thinning of the ozone layer is harmful to man. Explain why this is so.
(ii) Explain, with essential details, how the ozone layer is destroyed.
- (c) (i) Indicate clearly the organic compounds necessary for the production of each of the materials 'polyester' and 'nylon'.
N.B. You may use either words or structures for this purpose.
(ii) You are supplied a sample of polyester and a sample of nylon. Indicate clearly how you attempt to chemically distinguish between these two samples.
- (d) Propose **three** chemical tests that could be performed easily for the purpose of clearly demonstrating that P—H bonds are present in H_3PO_2 . You should also give the relevant observations along with the tests.

Clue: Think of oxidation numbers.