

**45th INTERNATIONAL
CHEMISTRY OLYMPIAD**

2013

UK Round One

STUDENT QUESTION BOOKLET

* * * * *

- The time allowed is 2 hours.
- Attempt all 5 questions.
- Write your answers in the special answer booklet.
- In your calculations, please write only the essential steps in the answer booklet.
- Always give the appropriate units and number of significant figures.
- You are provided with a copy of the Periodic Table.
- Do *NOT* write anything in the right hand margin of the answer booklet.
- The marks available for each question are shown below; this may be helpful when dividing your time between questions.

Question	1	2	3	4	5	Total
Marks Available	9	13	9	15	17	63

Some of the questions will contain material you will not be familiar with. However, by logically applying the skills you have learnt as a chemist, you should be able to work through the problems. There are different ways to approach the tasks – even if you cannot complete certain parts of a question, you may still find subsequent parts straightforward.

H 1 1.008																	He 2 4.003
Li 3 6.94																	Ne 10 20.18
Na 11 22.99																	Ar 18 39.95
K 19 39.102	Ca 20 40.08	Sc 21 44.96	Ti 22 47.90	V 23 50.94	Cr 24 52.00	Mn 25 54.94	Fe 26 55.85	Co 27 58.93	Ni 28 58.71	Cu 29 63.55	Zn 30 65.37	Ga 31 69.72	Ge 32 72.59	As 33 74.92	Se 34 78.96	Br 35 79.904	Kr 36 83.80
Rb 37 85.47	Sr 38 87.62	Y 39 88.91	Zr 40 91.22	Nb 41 92.91	Mo 42 95.94	Tc 43	Ru 44 101.07	Rh 45 102.91	Pd 46 106.4	Ag 47 107.87	Cd 48 112.40	In 49 114.82	Sn 50 118.69	Sb 51 121.75	Te 52 127.60	I 53 126.90	Xe 54 131.30
Cs 55 132.91	Ba 56 137.34	La* 57 138.91	Hf 72 178.49	Ta 73 180.95	W 74 183.85	Re 75 186.2	Os 76 190.2	Ir 77 192.2	Pt 78 195.09	Au 79 196.97	Hg 80 200.59	Tl 81 204.37	Pb 82 207.2	Bi 83 208.98	Po 84	At 85	Rn 86
Fr 87	Ra 88	Ac+ 89															

symbol
atomic number
relative atomic mass

*Lanthanides	Ce 58 140.12	Pr 59 140.91	Nd 60 144.24	Pm 61	Sm 62 150.4	Eu 63 151.96	Gd 64 157.25	Tb 65 158.93	Dy 66 162.50	Ho 67 164.93	Er 68 167.26	Tm 69 168.93	Yb 70 173.04	Lu 71 174.97
+Actinides	Th 90 232.01	Pa 91	U 92 238.03	Np 93	Pu 94	Am 95	Cm 96	Bk 97	Cf 98	Es 99	Fm 100	Md 101	No 102	Lr 103


1. This question is about rocket fuels

The Mars Curiosity rover's landing in August 2012 was achieved using variable thrust mono propellant hydrazine rocket thrusters. Hydrazine, N_2H_4 , is popular with NASA as it produces no carbon dioxide.


The hydrazine is passed over a suitable catalyst and decomposes to its elements. The rapid production of the hot gaseous elements is what provides the thrust. Ammonia can be formed as an intermediate during the decomposition.



- (a) Write a balanced equation for hydrazine decomposing to ammonia and nitrogen gas.
- (b) Hydrazine may be obtained from the reaction between ammonia and hydrogen peroxide.




Work out the standard enthalpy change for the decomposition of hydrazine to its elements. The standard enthalpy changes of formation in kJ mol^{-1} are:




- (c) The first ever rocket-powered fighter plane, the Messerschmitt Me 163, was powered by the reaction between a hydrazine-methanol mixture, known as 'C-Stoff', and hydrogen peroxide ('T-Stoff').

- (i) Hydrogen peroxide reacts with the hydrazine as shown in the equation.



State the oxidation number of nitrogen and oxygen in the reactants and products.

- (ii) Hydrogen peroxide oxidises the methanol to carbon dioxide and water. Write a balanced equation for this reaction.
- (iii) The fighter plane would hold 225 litres of hydrazine and 862 litres of methanol. Use the following standard enthalpy changes and densities to calculate the heat energy evolved under standard conditions for the combustion of this quantity of rocket fuel. Assume that all the hydrazine and methanol are fully combusted.



- (d) Hydrazine is also commonly combined with dinitrogen tetroxide, N_2O_4 , in rocket fuels. This forms a hypergolic mixture, i.e. the reactants ignite spontaneously on contact. NASA used $\text{N}_2\text{H}_4 / \text{N}_2\text{O}_4$ in many space vehicles and it is likely to be used in next-generation vehicles.
- (i) Reactions used in rocketry produce chemically stable products (making the reaction exothermic) that are formed as gases (which provide thrust). Suggest the reaction products that are formed in the reaction between N_2H_4 and N_2O_4 .
- (ii) Pure N_2O_4 , when warmed, initially decomposes not into its elements but instead forms a brown gas. Suggest the identity of this brown gas.
- (e) A derivative of hydrazine with formula $\text{C}_2\text{H}_8\text{N}_2$ was used in rocket fuels in the Apollo missions. It has two nitrogen atoms that are in different chemical environments and two carbon atoms that are in the same chemical environment. Draw the structure of $\text{C}_2\text{H}_8\text{N}_2$.



2. This question is about Great Britain being better than any other country at cycling

One of the most successful British sports at the London 2012 Olympics was cycling. British cyclists won eight gold, two silver and two bronze medals. In addition, Bradley Wiggins became the first ever Briton to win the Tour de France.



- (a) Bronze medals contain copper. A 0.800 g sample of a bronze medal was dissolved in hot concentrated nitric acid. After cooling and dilution, an excess of potassium iodide solution was added and the solution was made up to 250.0 cm³. A 25.00 cm³ aliquot of this solution required 12.20 cm³ of 0.100 mol dm⁻³ sodium thiosulfate solution in the presence of starch indicator.

Calculate the percentage by mass of copper in the bronze medal.



- (b) All medals that were won at London 2012 had a diameter of 85 mm and were 7 mm thick. The silver medal was made up of 92.5% silver and 7.5% copper, by mass.

Calculate the mass of a silver medal. Assume that the density of the alloy varies proportionally to its composition by mass.

Densities in g cm⁻³: Ag, 10.49; Cu, 8.96

- (c) Gold medals are made of gold, silver and copper. A 5.000 g sample of a gold medal was warmed with excess concentrated nitric acid. An undissolved residue was separated by filtration, washed, dried and weighed. Its mass was 0.067 g. Then an excess of dilute hydrochloric acid was added to the solution in nitric acid. The precipitate formed was separated by filtration, washed, dried and weighed. Its mass was 6.144 g.

Calculate the percentage by mass of gold, silver and copper in the gold medal.

The GB cyclists have individually designed bikes, which have to cope with the demands placed on them whilst being as light as possible. One way a light bike could be achieved is by altering the gas used to inflate the bike tyres.

You may assume that for the remainder of this question that gases behave as ideal gases and that they follow the ideal gas law:

$$pV = nRT$$


where p = pressure in Pa
 V = volume in m^3
 n = number of moles
 R (the gas constant) = $8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
 T = temperature in Kelvin

The tyres on a bike are approximately the shape of a torus. The equation for the volume of a torus, V , is given below.

$$V = \frac{\pi^2 r d^2}{2}$$

r = distance from centre of torus to **centre** of tyre tube

d = diameter of tyre tube



(d) A typical bike may have a wheel diameter of 66 cm and a tyre diameter of 23 mm.

Calculate the volume of a tyre, in m^3 .

If you are unable to calculate the volume of a tyre, you may use the value of 0.001 m^3 in the subsequent parts of this question.

(e) (i) The air pressure used in the tyres is typically 120 psi, much higher than atmospheric pressure.

$$1 \text{ psi} = 6895 \text{ Pa.}$$

Calculate the number of moles of gas in a tyre, at $25 \text{ }^\circ\text{C}$.

(ii) Assuming air is a mixture of 80 % nitrogen gas and 20 % oxygen gas, calculate the total mass of air in **both** tyres on a bike.

(iii) In cycling the smallest of changes can make the difference between winning and losing. The small reduction in mass upon inflating tyres with helium instead of air would be worth considering if it was not for that fact that the very small helium atoms escape through the rubber of tyres much more rapidly.

Calculate the reduction in mass of the bike if both tyres were inflated with helium instead of air.


(iv) SF_6 is one of the densest substances that would still remain in the gas phase at this pressure.

What would be the increase in mass if the bike tyres were filled with SF_6 ?

3. This question is about chemistry general knowledge based on coloured compounds









Chemists recognise many substances by their colour.

This question tests your knowledge about the colour of a range of chemicals that you will have come across as part of your studies of chemistry.



- (a) In the answer booklet, a number of colours are listed. For each colour, if one of the substances below is that colour, give its letter. If that colour cannot be made using a single substance, give the letters of two substances that will produce this colour when mixed. A substance may be used more than once.

Choose from



- (b) A student was provided with 7 unknown aqueous solutions. Each solution contained only one substance and all solutions were different.

The student mixed pairs of solutions in an attempt to identify each of the substances. The results from these tests are shown below. A blank cell shows where no observable results were obtained.

Analyse the results and assign the following substances as solutions **T-Z**.

barium chloride

dissolved chlorine gas
(i.e. chlorine water)

iron(II) sulfate

lead(II) nitrate

silver nitrate

sodium carbonate

sodium iodide


	T	U	V	W	X	Y	Z
T		Bright yellow precipitate formed	White precipitate formed	–	White precipitate formed	White precipitate formed	White precipitate formed
U	Bright yellow precipitate formed		–	Yellow precipitate formed	–	–	Brown solution
V	White precipitate formed	–		White precipitate formed (darkened in daylight)	White precipitate formed	White precipitate formed	–
W	–	Yellow precipitate formed	White precipitate formed (darkened in daylight)		Off-white precipitate formed	White precipitate formed	White precipitate formed (darkened in daylight)
X	White precipitate formed	–	White precipitate formed	Off-white precipitate formed		Dirty green precipitate formed (turned brown on standing)	–
Y	White precipitate formed	–	White precipitate formed	White precipitate formed	Dirty green precipitate formed (turned brown on standing)		Pale yellow solution
Z	White precipitate formed	Brown solution	–	White precipitate formed (darkened in daylight)	–	Pale yellow solution	

4. This question is about GPs giving out too many drugs

The drugs diazepam (Valium®) and alprazolam (Xanax®) belong to a family of compounds called benzodiazepines which were once hailed as the answer to many ailments. In 1988, The Committee on Safety of Medicines advised GPs to limit their prescriptions of these drugs due to their overuse; however, in England in 2010 more than 6.6 million benzodiazepine prescriptions for anxiety were dispensed. Heath Ledger, Michael Jackson and Whitney Houston are among some of the many people that are believed to have died from drug overdoses that involved benzodiazepines.



The synthesis of diazepam from 4-chloroaniline is shown below. The percentage yield for each step is also shown. By-products are not always shown.




(a) (i) Calculate the overall percentage yield for the conversion of 4-chloroaniline to diazepam.

(ii) A patient was prescribed diazepam for three years at a dose of 5 mg, four times a day.


Calculate the mass of diazepam this was, and hence the mass of 4-chloroaniline needed to make the drug for this patient.

(b) Draw the structures of intermediates **A** – **H** in the synthesis of diazepam.

In the liver it was found that diazepam underwent a demethylation reaction. The product of this demethylation reaction was used as inspiration for the synthesis of the drug alprazolam.




(c) The synthesis of alprazolam is shown below. Draw the structures of intermediates **I**, **J** and **K**



(d) In the final stage of the synthesis, compound **L** is treated with a chemical called 'DEAD' to convert it into alprazolam. DEAD stands for diethyl azodicarboxylate.

How would you classify the reaction of **L** to alprazolam using one of the terms below?

Tick the correct answer in your answer booklet.



Isomerisation

Hydrolysis


Condensation

Oxidation

Reduction

5. This question is about getting big muscles

Creatine has recently become one of the most widely used nutritional supplements among athletes. Although there is much debate about which of the advertised beneficial effects of creatine are actually true, the use of creatine is generally believed to lead to a short-term gain in body mass/muscle size. The structure of creatine is shown below.







In the body, creatine is converted into phosphocreatine which is used as an energy reserve in the muscles that can be rapidly mobilised to convert adenosine diphosphate (ADP) back into adenosine triphosphate (ATP) – the body's energy currency – in times of need.



- (a) Creatine is often sold in capsules labelled as 'Pure Creatine Monohydrate'. Write the molecular formula of creatine monohydrate.
- (b) Creatine is synthesised naturally in organisms from three amino acids: glycine, methionine and arginine.

In your answer booklet, for each of the carbon atoms in creatine (labelled as **1-4**), suggest which carbon atom of the three amino acids it came from (labelled as **A-M**)



- (c) Like amino acids, creatine exists in different ionised forms depending on the pH of the solution it is in. This causes the overall charge on the molecule to vary. Draw the most common form of creatine at each of the following pHs (the overall charge on the molecule at each pH is given).
- (i) pH 1 (overall charge = + 1) (ii) pH 7 (overall charge = neutral) (iii) pH 12 (overall charge = - 1)

The chemical structure of creatine and these amino acids can be analysed by ¹H NMR.

As these are polar molecules, the NMR spectra are run in D₂O solvent. In D₂O, protons attached to nitrogen or oxygen atoms undergo rapid exchange with deuteriums from the solvent. This means that by the time the NMR is run, all N-H bonds have been replaced by N-D bonds and all O-H bonds by O-D bonds. As signals from deuterium atoms are not observed in ¹H NMR spectra, no signals from N-H or O-H groups in the molecule are seen in the spectrum.

The number of signals observed depends on the symmetry of the molecule. Each hydrogen atom in a unique environment gives rise to a signal at a different chemical shift in the spectrum. Occasionally, signals from two different environments can appear on top of one another when the difference in chemical shifts between the environments is very small.


The area under each signal is proportional to the number of protons in that environment. This is shown by an integral trace (the stepped line on the spectrum). The height of each step is proportional to the area under that signal.

The appearance of the signals can be complicated by coupling. If the hydrogen atom(s) are within three bonds of another hydrogen which is in a different environment, instead of appearing as a single peak, its signal is split into a number of peaks. If the hydrogen under consideration is within three bonds of **n** hydrogens in a different environment from itself, it will be split into **(n + 1) equally spaced** peaks. The ratio of the area under the peaks is given by the number in Pascal's triangle (shown on the right). Due to rapid exchange of any protons/deuteriums bonded to oxygen or nitrogen atoms with the solvent, no coupling is seen to protons/deuteriums bonded to oxygen or nitrogen atoms.

n	Intensities of peaks
0	1
1	1 : 1
2	1 : 2 : 1
3	1 : 3 : 3 : 1
4	1 : 4 : 6 : 4 : 1

(d) Consider the amino acid methionine.

Complete the table in the answer booklet for carbons **C**, **D** and **F** in methionine to suggest the appearance of the overall signal from the protons bonded to that carbon atom.



(e) Usually all protons attached to the same carbon atom are in the same chemical environment; however, this is not always the case. Two protons on the same carbon atom that are in different chemical environments are called diastereotopic protons.

These are most often observed where the carbon under consideration is bonded to an asymmetric carbon atom. An asymmetric carbon atom has four different chemical groups attached to it.


Consider glycine, methionine and arginine. In these three amino acids, write the letters of all such carbons whose diastereotopic protons would be observed as different signals in their spectra.

In the body, creatine is in equilibrium with a cyclic molecule called creatinine, by the following equation. The position of equilibrium varies with pH.



Creatinine is a metabolic waste product that is not used by the body. It is filtered out in the kidneys.

The ^1H NMR spectrum in D_2O of a creatine/creatinine solution is shown below. Three signals are observed. Creatinine gives rise to signal A. Creatine gives rise to signal B. Both creatine and creatinine give rise to signal C.




- (f) Suggest a structure for creatinine.
- (g) Assuming this sample has reached equilibrium, calculate a value for the equilibrium constant, K , at this pH and temperature. Show clearly how you worked this out. You may ignore the concentration of water in your calculation.
- (h) A problem with creatine supplementation is that a lot of the creatine taken does not get absorbed by the body. Recently, supplements containing derivatives of creatine have been marketed. These are usually more lipophilic (dissolve more easily in fats) in an effort to improve uptake into the body.

The ^1H NMR spectrum in D_2O of one of these supplements is shown below. Some regions of the spectrum have been expanded on the left hand side of the figure to help with your analysis.

This supplement exists in an ionised form at pH 1 but does not exist in an ionised form at pH 12.

Suggest a structure for this supplement.



Acknowledgements & References

Q1

The Curiosity Mars rover image is courtesy of NASA.

Q2

Information on the composition of the medals was taken from www.london2012.com.

The Bradley Wiggins image is courtesy of Getty Images.

Q3

The image of the test tubes is courtesy of iStockphoto.

Q4

The Independent **Thursday 29th December 2011** Doctors sued for creating 'Valium addicts' *Nina Lakhani*

The image of diazepam is courtesy of Paul Brown and Alamy.

Heterocycles **1981** 16 1491-1494

Journal of Medicinal Chemistry **1977** 20 1694-1697

Journal of Medicinal Chemistry **1979** 22 1-7

Journal of Organic Chemistry **1980** 45 1675-1681

Tetrahedron Letters **1971** 12 1609-1612

Q5

John Gallacher is thanked for providing an image of his arm.

The University of Oxford is thanked for use of their NMR facilities.


The creatine supplements analysed were from Precision Engineered Ltd.

Biochemical Journal **1928** 22 920-929

Journal of the Chemical Society, Perkin Transactions II **1985** 1465-1467

The UK Chemistry Olympiad is supported by INEOS. INEOS is a leading global manufacturer of petrochemicals, specialty chemicals and oil products. It comprises 19 businesses each with a major chemical company heritage. The production network spans 73 manufacturing facilities in 19 countries throughout the world. The chemicals INEOS produce enhance almost every aspect of modern life.

INEOS
THE WORD FOR CHEMICALS



45th INTERNATIONAL CHEMISTRY OLYMPIAD

UK Round 1 - 2013

MARK SCHEME

Question	1	2	3	4	5	Total
Mark	9	13	9	15	17	63

Although we would encourage students to always quote answers to an appropriate number of significant figures, do not penalise students for significant figure errors. Allow where a student's answers differ slightly from the mark scheme due to the use of rounded/non-rounded data from an earlier part of the question.

For answers with missing or incorrect units, penalise 1 mark for the first occurrence in **each** question. Do not penalise for subsequent occurrences in the same question.

Question 1																		
(a)		$3\text{N}_2\text{H}_4(\text{l}) \rightarrow 4\text{NH}_3(\text{g}) + \text{N}_2(\text{g})$ [Ignore state symbols]	1															
(b)		$\Delta_f H^\ominus = ((2 \times 46.1) + 187.8 - (2 \times 285.8) + \Delta_f H^\ominus (\text{N}_2\text{H}_4)) \text{ kJ mol}^{-1} = -241.0 \text{ kJ mol}^{-1}$ $\Delta_f H^\ominus (\text{N}_2\text{H}_4) = 50.6 \text{ kJ mol}^{-1}$ Decomposition enthalpy = $-50.6 \text{ kJ mol}^{-1}$ [Positive answers of correct magnitude do not score credit.]	1															
(c)	(i)	<table border="1"> <thead> <tr> <th></th> <th>N_2H_4</th> <th>H_2O_2</th> <th>N_2</th> <th>H_2O</th> </tr> </thead> <tbody> <tr> <td>Ox. state of N</td> <td>-2</td> <td></td> <td>0</td> <td></td> </tr> <tr> <td>Ox. state of O</td> <td></td> <td>-1</td> <td></td> <td>-2</td> </tr> </tbody> </table> [No partial credit given]		N_2H_4	H_2O_2	N_2	H_2O	Ox. state of N	-2		0		Ox. state of O		-1		-2	1
	N_2H_4	H_2O_2	N_2	H_2O														
Ox. state of N	-2		0															
Ox. state of O		-1		-2														

	(ii)	$\text{CH}_3\text{OH}(\text{l}) + 3\text{H}_2\text{O}_2(\text{l}) \rightarrow \text{CO}_2(\text{g}) + 5\text{H}_2\text{O}(\text{l})$ [Ignore state symbols]	1
	(iii)	Amount of hydrazine = $225000 \text{ cm}^3 \times 1.021 \text{ g cm}^{-3} / 32.052 \text{ g mol}^{-1}$ = 7167 mol Amount of methanol = $862000 \text{ cm}^3 \times 0.7918 \text{ g cm}^{-3} / 32.042 \text{ g mol}^{-1}$ = 21301 mol [Both amounts needed for one mark] Heat energy evolved from hydrazine = $7167 \text{ mol} \times 622.2 \text{ kJ mol}^{-1}$ = $4.459 \times 10^6 \text{ kJ}$ Heat energy evolved from methanol = $21301 \text{ mol} \times 726.0 \text{ kJ mol}^{-1}$ = $15.465 \times 10^6 \text{ kJ}$ Total heat energy evolved from oxidation of rocket fuel = $19.9 \times 10^6 \text{ kJ}$ [Correct answer scores both marks. Accept $-19.9 \times 10^6 \text{ kJ}$]	1 1
(d)	(i)	N_2 and H_2O [Half a mark each. Accept 'nitrogen and water'.]	1
	(ii)	NO_2 [Accept 'nitrogen dioxide'.]	1
(e)		$(\text{CH}_3)_2\text{N}-\text{NH}_2$ <i>This is known in the trade as UDMH (unsymmetrical dimethylhydrazine)</i>	1
Total			9

Question 2

(a)	Amount of $\text{S}_2\text{O}_3^{2-} = 0.0122 \text{ dm}^3 \times 0.100 \text{ mol dm}^{-3} = 1.22 \times 10^{-3} \text{ mol}$ Amount of Cu = $1.22 \times 10^{-2} \text{ mol}$ Mass of Cu = $1.22 \times 10^{-2} \text{ mol} \times 63.55 \text{ g mol}^{-1} = 0.775 \text{ g}$ Percentage of Cu by mass = $100 \% \times 0.775 \text{ g} / 0.800 \text{ g} = 96.9 \%$	1
(b)	Volume of medal = $\pi r^2 h = \pi \times (4.25 \text{ cm})^2 \times 0.7 \text{ cm} = 39.72 \text{ cm}^3$ Density of medal = $(0.925 \times 10.49 \text{ g cm}^{-3}) + (0.075 \times 8.96 \text{ g cm}^{-3}) = 10.38 \text{ g cm}^{-3}$ Mass of medal = $39.72 \text{ cm}^3 \times 10.38 \text{ g cm}^{-3} = 412 \text{ g}$ [Correct answer scores both marks.]	1 1

(c)	<p>Mass of Au = 0.067 g</p> <p>Amount of Ag = amount of AgCl = $6.144 \text{ g} / (107.87 + 35.45) \text{ g mol}^{-1} = 4.287 \times 10^{-2} \text{ mol}$</p> <p>Mass of Ag = $4.287 \times 10^{-2} \text{ mol} \times 107.87 \text{ g mol}^{-1} = 4.624 \text{ g}$</p> <p>Mass of Cu = $5.000 \text{ g} - 0.067 \text{ g} - 4.624 \text{ g} = 0.309 \text{ g}$</p> <p>Percentage of Au by mass = $100 \% \times 0.067 \text{ g} / 5.000 \text{ g} = 1.34 \%$</p> <p>Percentage of Ag by mass = $100 \% \times 4.624 \text{ g} / 5.000 \text{ g} = 92.5 \%$</p> <p>Percentage of Cu by mass = $100 \% \times 0.309 \text{ g} / 5.000 \text{ g} = 6.18 \%$</p> <p>[One mark awarded for each correct percentage. Allow error carried forward in the copper percentage. Allow minor differences due to rounding.]</p>	<p>1</p> <p>1</p> <p>1</p>
(d)	<p>$d = \text{tyre diameter} = 0.023 \text{ m}$</p> <p>$r = (\text{wheel diameter} / 2) - (\text{tyre diameter} / 2) = 0.33 \text{ m} - 0.0115 \text{ m}$ $= 0.3185 \text{ m}$</p> <p>[One mark for correct value of r]</p> <p>volume = $\pi^2 \times 0.3185 \text{ m} \times (0.023 \text{ m})^2 / 2$ $= 8.314 \times 10^{-4} \text{ m}^3$</p> <p>[Correct answer scores both marks.]</p>	<p>1</p> <p>1</p>
(e)	<p>(i) $p = 8.27 \times 10^5 \text{ Pa}$; $V = 8.31 \times 10^{-4} \text{ m}^3$; $T = 298 \text{ K}$</p> <p>$n = pV/RT$</p> <p>[One mark for correct method.]</p> <p>$n = (8.27 \times 10^5 \text{ Pa} \times 8.31 \times 10^{-4} \text{ m}^3) / (8.314 \text{ J K}^{-1} \text{ mol}^{-1} \times 298 \text{ K})$ $n = 0.278 \text{ mol}$</p> <p>[Correct answer scores both marks; $n = 0.334 \text{ mol}$ if value of 0.001 m^3 used for volume.]</p>	<p>1</p> <p>1</p>
	<p>(ii) $N_2 = 28.02 \text{ g mol}^{-1}$; $O_2 = 32.00 \text{ g mol}^{-1}$</p> <p>mass in one tyre = $((0.8 \times 28.02 \text{ g mol}^{-1}) + (0.2 \times 32.00 \text{ g mol}^{-1})) \times 0.278 \text{ mol}$ mass in one tyre = 8.011 g</p> <p>mass of air in both tyres = $8.011 \text{ g} \times 2$ $= 16.02 \text{ g}$</p> <p>[Mass = 19.25 g if value of 0.001 m^3 used for volume. Allow any approximations that are more accurate than this, for example if the student has decided to use 78% N_2, 21% O_2, 1% Ar.]</p>	<p>1</p>
	<p>(iii) $He = 4.003 \text{ g mol}^{-1}$</p> <p>mass = $2 \times 0.278 \text{ mol} \times 4.003 \text{ g mol}^{-1}$ mass = 2.226 g</p> <p>mass reduction = $16.02 \text{ g} - 2.226 \text{ g}$ mass reduction = 13.79 g</p> <p>[Error carried forward: accept answer from (e)(ii) minus 2.226 g or answer from (e)(ii) minus 2.674 g if 0.001 m^3 used for volume.]</p>	<p>1</p>




	<p><i>Although this mass reduction is small, it is significant enough to be considered. Unfortunately being very small, helium escapes through the rubber of tyres much more easily and so is rarely used.</i></p>	
(iv)	$SF_6 = 32.06 \text{ g mol}^{-1} + (6 \times 19.00 \text{ g mol}^{-1}) = 146.06 \text{ g mol}^{-1}$ $\text{mass} = 2 \times 0.278 \text{ mol} \times 146.06 \text{ g mol}^{-1}$ $\text{mass} = 81.209 \text{ g}$ $\text{mass increase} = 81.209 \text{ g} - 16.02 \text{ g}$ $\text{mass increase} = 65.19 \text{ g}$ <p>[Error carried forward: accept 81.209 g minus answer from (e)(ii), or 97.568 g minus answer from (e)(ii) if 0.001 m³ used for volume.]</p>	1
Total		13


Question 3

(a)	<table border="1"> <tr><td>Red</td><td>C</td><td></td></tr> <tr><td>Orange</td><td>G</td><td></td></tr> <tr><td>Yellow</td><td>G</td><td>H</td></tr> <tr><td>Green</td><td>F</td><td></td></tr> <tr><td>Blue</td><td>D</td><td>H</td></tr> <tr><td>Violet</td><td>I</td><td></td></tr> </table> <p>[Award half a mark for each colour. For yellow and blue both letters are needed to score the half mark for that colour.]</p>	Red	C		Orange	G		Yellow	G	H	Green	F		Blue	D	H	Violet	I		3
Red	C																			
Orange	G																			
Yellow	G	H																		
Green	F																			
Blue	D	H																		
Violet	I																			
(b)	<table border="1"> <tr><td>T</td><td>lead(II) nitrate</td></tr> <tr><td>U</td><td>sodium iodide</td></tr> <tr><td>V</td><td>barium chloride</td></tr> <tr><td>W</td><td>silver nitrate</td></tr> <tr><td>X</td><td>sodium carbonate</td></tr> <tr><td>Y</td><td>iron(II) sulfate</td></tr> <tr><td>Z</td><td>chlorine water/dissolved chlorine gas</td></tr> </table> <p>All 7 correct = 6 marks 5 or 6 correct = 5 marks 4 correct = 4 marks 3 correct = 3 marks 2 correct = 2 marks 1 correct = 1 mark</p> <p>[Ignore spelling errors as long as substance is recognisable. Oxidation states not needed. Accept if correct chemical formulae have been written instead of words.]</p>	T	lead(II) nitrate	U	sodium iodide	V	barium chloride	W	silver nitrate	X	sodium carbonate	Y	iron(II) sulfate	Z	chlorine water/dissolved chlorine gas	6				
T	lead(II) nitrate																			
U	sodium iodide																			
V	barium chloride																			
W	silver nitrate																			
X	sodium carbonate																			
Y	iron(II) sulfate																			
Z	chlorine water/dissolved chlorine gas																			
Total		9																		


Question 4

(a)	(i)	10.8 %	1
<p><i>More modern syntheses have considerably improved upon this overall yield.</i></p>			


<p>(ii)</p>	<p>Mass of diazepam per dose = 5×10^{-3} g</p> <p>Total mass of diazepam = 5×10^{-3} g \times 4 \times 365 \times 3 Total mass of diazepam = 21.9 g [Allow an extra day added for a leap year.]</p> <p>Amount of diazepam = 21.9 g / 284.734 g mol⁻¹ Amount of diazepam = 0.0769 mol</p> <p>Amount of 4-chloroaniline = 0.0769 mol / 0.108 Amount of 4-chloroaniline = 0.712 mol Molecular formula of 4-chloroaniline = C₆H₆NCl M_r of 4-chloroaniline = (6 \times 12.01 g mol⁻¹) + (6 \times 1.008 g mol⁻¹) + 35.45 g mol⁻¹ + 14.01 g mol⁻¹ M_r of 4-chloroaniline = 127.568 g mol⁻¹</p> <p>Mass of 4-chloroaniline = 0.712 mol \times 127.568 g mol⁻¹ Mass of 4-chloroaniline = 90.8 g</p> <p>[Correct answer scores both marks. Error carried forward: accept answers based on incorrect answer to (a)(i)]</p>	<p>1</p> <p>1</p>
<p>(b)</p>	<p>Structure of A</p>  <p>[If bromine atom is in wrong position on benzene ring, no credit is given here, but full credit is awarded in B, D, E and F provided rest of structure correct.]</p>	<p>1</p>
	<p>Structure of B</p> 	<p>1</p>
	<p>Structure of C</p> 	<p>1</p>

Structure of **D**


1

Structure of **E**


1

Structure of **F**

1




Structure of **G**

1

Structure of **H**





1

[Allow stereoisomer with other geometry around C=N bond.]

(c)	<p style="text-align: center;">Structure of I</p> 	1										
	<p style="text-align: center;">Structure of J</p>  <p>[Accept either structure. Also allow stereoisomer with other geometry around the top C=N bond in the right hand structure.] <i>The left structure is believed to be the predominant one.</i></p>	1										
	<p style="text-align: center;">Structure of K</p>  <p>[Accept either structure and also any stereoisomers with alternative geometries around C=N bonds not in the ring.] <i>The left structure is believed to be the predominant one.</i></p>	1										
(d)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 20%;">Isomerisation</th> <th style="width: 20%;">Hydrolysis</th> <th style="width: 20%;">Condensation</th> <th style="width: 20%;">Oxidation</th> <th style="width: 20%;">Reduction</th> </tr> </thead> <tbody> <tr> <td style="height: 20px;"></td> <td></td> <td></td> <td style="text-align: center;">✓</td> <td></td> </tr> </tbody> </table> <p><i>DEAD is also sometimes known as DEADCAT.</i></p>	Isomerisation	Hydrolysis	Condensation	Oxidation	Reduction				✓		1
Isomerisation	Hydrolysis	Condensation	Oxidation	Reduction								
			✓									
Total		15										

Question 5

(a)	<p>$C_4H_9N_3O_2 \cdot H_2O$ or $C_4H_{11}N_3O_3$ [Accept answers where order of elements is different.]</p>	1
-----	---	---

(b)		<table border="1" data-bbox="616 152 1031 367"> <thead> <tr> <th>Carbon Atom in Creatine</th> <th>Carbon Atom in Amino Acid</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>H</td> </tr> <tr> <td>2</td> <td>C</td> </tr> <tr> <td>3</td> <td>A</td> </tr> <tr> <td>4</td> <td>B</td> </tr> </tbody> </table> <p data-bbox="233 398 1286 430">All four correct = 3 marks Three correct = 2 marks Two correct = 1 mark</p>	Carbon Atom in Creatine	Carbon Atom in Amino Acid	1	H	2	C	3	A	4	B	3																		
Carbon Atom in Creatine	Carbon Atom in Amino Acid																														
1	H																														
2	C																														
3	A																														
4	B																														
(c)	(i)	<div style="display: flex; justify-content: space-around; align-items: center;">  </div> <p data-bbox="233 672 1391 770">[Any one of the three resonance structures shown above is worth 2 marks i.e. the extra proton must be on the uppermost nitrogen as drawn in the diagram for 2 marks. Either of the two structures shown below should be awarded 1 mark.]</p> <p data-bbox="233 770 1382 837"><i>The uppermost nitrogen is protonated as this maintains the delocalisation present in the neutral guanidinium group in the conjugate base form.</i></p> <div style="display: flex; justify-content: space-around; align-items: center;">  </div>	2																												
	(ii)	<div style="display: flex; justify-content: space-around; align-items: center;">  </div> <p data-bbox="233 1281 1398 1415">[The first structure is chemically correct, but any one of the three structures shown above is worth 1 mark. – The mark for protonating the correct nitrogen is awarded in part (i). Allow the other two resonance forms for the guanidinium group in the structure on the left as in part (i)]</p>	1																												
	(iii)	<div style="display: flex; justify-content: center; align-items: center;">  </div>	1																												
(d)		<table border="1" data-bbox="248 1693 1366 1966"> <thead> <tr> <th>Carbon atom</th> <th>Singlet</th> <th>1:1 doublet</th> <th>1:2:1 triplet</th> <th>1:3:3:1 quartet</th> <th>1:4:6:4:1 quintet</th> <th>No signal observed</th> </tr> </thead> <tbody> <tr> <td>C</td> <td style="text-align: center;">✓</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>D</td> <td></td> <td></td> <td style="text-align: center;">✓</td> <td></td> <td></td> <td></td> </tr> <tr> <td>F</td> <td></td> <td></td> <td style="text-align: center;">✓</td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	Carbon atom	Singlet	1:1 doublet	1:2:1 triplet	1:3:3:1 quartet	1:4:6:4:1 quintet	No signal observed	C	✓						D			✓				F			✓				1 1 1
Carbon atom	Singlet	1:1 doublet	1:2:1 triplet	1:3:3:1 quartet	1:4:6:4:1 quintet	No signal observed																									
C	✓																														
D			✓																												
F			✓																												

(e)	<p>E and K</p> <p>[Award half a mark each. If other letters are written minus half a mark for each other letter down to zero.]</p>	1
(f)	<p>[Any one of the five alternatives below is to be awarded the mark.] <i>The percentage of each tautomer is solvent dependent, although the top two are by far the most important. In protic solvents, hydrogen-bonding favours the top left structure.</i></p> <div style="text-align: center;"> </div>	1
(g)	<p>$K = [\text{Creatinine}] / [\text{Creatine}]$ $K = \text{Integral height of signal A} / \text{Integral height of signal B}$ $K = 4$</p> <p>[This has no units. Award values between 3.5 and 5.0 the mark. There must be evidence of working/using the correct integral method to gain the mark.] <i>Creatinine is favoured at more acidic pHs and creatine at more alkaline pHs.</i></p>	1
(h)	<div style="text-align: center;"> </div> <p>[The correct structure is to be awarded 3 marks. The hydrochloride salt of this molecule (protonated on any one nitrogen) should also be awarded 3 marks. Incorrect structures may score 2 marks if they obey any two of the three criteria below, and 1 mark for obeying any one of the criteria.]</p> <ul style="list-style-type: none"> ■ A total of 10 C–H protons in the molecule. <i>This shows the student has successfully used the integrals in the spectrum to calculate the number of hydrogens.</i> ■ The presence of a discrete ethyl group in the molecule. <i>This shows the student has understood the coupling patterns in the NMR.</i> ■ The presence of an ester functional group in place of the carboxylic acid. <i>This shows the student has understood the ionisation states of the molecule at different pHs.</i> 	3
Total		17